




**You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice**

Title: *Macrosiphoniella pulvera* (Walker, 1848) an aphid species (Hemiptera: Aphididae) new to the Polish fauna

Author: Natalia Kaszyca-Taszakowska, Łukasz Depa

Citation style: Kaszyca-Taszakowska Natalia, Depa Łukasz. (2019). *Macrosiphoniella pulvera* (Walker, 1848) an aphid species (Hemiptera: Aphididae) new to the Polish fauna. "Annals of the Upper Silesian Museum in Bytom, Entomology" Vol. 28 (2019), s. 1-3, doi 10.5281/zenodo.3557243



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

NATALIA KASZYCA-TASZAKOWSKA^{1*} , ŁUKASZ DEPA¹ 

***Macrosiphoniella pulvera* (WALKER, 1848) an aphid species (Hemiptera: Aphididae) new to the Polish fauna**

<http://doi.org/10.5281/zenodo.3557243>

¹ Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, Bankowa 9, 40-007 Katowice, Poland

* e-mail: nkaszyca@us.edu.pl

Abstract: The paper presents data on the occurrence of aphid species *Macrosiphoniella pulvera* in Poland.

Key words: Sternorrhyncha, new record, biodiversity, Poland.

INTRODUCTION

Aphids, according to most of other insects, are a group reaching its highest species diversity in the zone of moderate climates. Poland is among the best studied European countries in terms of aphid fauna, with 768 recorded aphid taxa (WOJCIECHOWSKI *et al.* 2015, KANTURSKI *et al.* 2017, KANTURSKI *et al.* 2018, MIŚKOWIEC *et al.* 2019). There are about 110 species of *Macrosiphoniella* worldwide, and only 21 species in Poland (OSIADACZ & HAŁAJ 2009). There is no host alternation and all species feed on daisies (Asteraceae), most on species of the division Anthemideae. Many *Macrosiphoniella* species feed on just one genus of this group, some on just one species. *Macrosiphonella pulvera* it feeds on several species of *Artemisia* plants: *Artemisia abrotanum*, *A. absinthium*, *A. adamsii*, *A. annua*, *A. austriaca*, *A. caerulea*, *A. herba-alba*, *A. mongolica*, *A. pontica*, *A. santonicum*, *A. serotina*, *A. sieversiana*, *A. szowitziana*, *A. terrae-albae*, *A. vulgaris* (BLACKMAN & EASTOP 2019). Here we present a record of aphid species of the genus *Macrosiphoniella* feeding on *Artemisia* new in Polish fauna.

MATERIAL AND METHODS

The species was collected during aphidological research conducted in the Silesian agglomeration. The specimens were found in the Kościuszko Park in Katowice (Upper Silesian Upland) (Fig.1.) (KONDRACKI 2013), on *Artemisia vulgaris*, on 20th June 2018. Aphid specimens were collected during the detailed searching of their host plants in field, preserved in 70% ethanol and mounted on microscopic slide for determination. After mounting on microscopic slide with method described by KANTURSKI & WIECZOREK (2012) the specimens were identified using online key by BLACKMAN & EASTOP (2019). The slides are collected in the entomological collection of the University of Silesia (DZUS).




Fig. 1. Map of species occurrence in Europe. The light green color indicates the countries in which *M. pulvera* occurs, dark green – the states in which its occurrence requires confirmation. The place where the aphid was found in Poland was marked with a red dot.

RESULTS AND DISCUSSION

Collection site: Katowice (UTM: CA56), roadside: 50°14'44.8"N 19°00'16.8"E, leg. Natalia Kaszyca-Taszakowska, det. Łukasz Depa; a single alate and aptera female (Fig. 2) and a few larvae were collected from *Artemisia vulgaris*. Greyish green wax-powdered with antennae pale with dark apices or dark from the middle of seg. III; legs brownish with darker tips of femora and tibiae. Siphunculi and cauda somewhat darker. Ant. seg. III with 11-23 rhinaria. The relatively low number of secondary rhinaria on antennal segment III and setae on this segment at most as long as its basal diameter are diagnostic features of this species, distinguishing it from very similar *M. abrotani*.

Widely distributed mostly in coastal regions of Europe and eastward across Asia to Mongolia; the Mongolian aphids, collected on *A. adamsi*, differed in some respects from European populations (HOLMAN & SZELEGIEWICZ 1978). This species is an ideal example of cryptic coloration due to their green color. The white wax powder on the *Macrosiphoniella pulvera* has the appearance of the cuticle surface of the *Artemisia* plants. This record establishes the total number of aphid taxa of specific and subspecific level in Poland to 769.

ACKNOWLEDGEMENTS

We wish to thank Mariusz Kanturski for verification of the identification.



Fig. 2. Alate female of *Macrosiphonella pulvera* – body ventral view and third antenna segment.

REFERENCES

- DURANTE M.P.M., EASTOP V., RAKAUSKAS R., REMAUDIERE G., NAFRIA J.M.N., HEIE O. 2013. Fauna Europaea: *Macrosiphoniella (Macrosiphoniella) pulvera* (WALKER, 1848). Fauna Europaea. Available at <http://faunaeur.org> (version: May 2019).
- BLACKMAN L., EASTOP F. 2019. Aphids on the World's Plants. An online information and information guide, <http://www.aphidsonworldsplants.info> (accessed 07.05.19).
- HOLMAN A., SZELEGIEWICZ H. 1978. Further aphids of the genus *Macrosiphoniella* (Homoptera, Aphididae) from Mongolia. *Acta Entomologica Bohemoslovaca* 75: 178–193.
- KANTURSKI M., LEE Y., DEPA Ł. 2018. New records of an alien aphid species *Tinocallis (Sappocallis) takachihoensis* from countries in central and northern Europe (Hemiptera, Aphididae, Calaphidinae). *ZooKeys* 730: 1–17.
- KANTURSKI M., MRUK K., MORAWSKI M., WOJCIECHOWSKI W., DEPA Ł. 2017. *Nearctaphis bakeri* (COWEN, 1895) and *Illinoia liriodendri* (MONELL, 1879) – two aphid species (Hemiptera: Aphididae) of alien origin new to the Polish fauna. *Annals of the Upper Silesian Museum in Bytom, Entomology* 26 (online 002): 1–6.
- KANTURSKI M., WIECZOREK K. 2012. Metody zbioru i preparowania mszyc (Hemiptera, Aphidoidea) w badaniach faunistycznych, taksonomicznych i molekularnych. *Młodzi Naukowcy dla polskiej Nauki* 8(5): 137–143.
- KONDRACKI J. 2013. Geografia regionalna Polski. Wydawnictwo Naukowe PWN. Warszawa: 444 pp.
- MIŚKOWIEC S., DEPA Ł., WĘGIEREK P., KASZYCA-TASZAKOWSKA N. 2019. Mszycy (Hemiptera, Aphidomorpha) rezerwatu przyrody „Dolina Żabnika”. *Acta entomologica silesiana* 27 (online 024): 1–9.
- OSIADACZ B., HALAJ R. 2009. The aphids (Hemiptera: Sternorrhyncha: Aphidinea) of Poland. A distributional checklist. *Polish Entomological Monographs* 6: 1–96.
- WOJCIECHOWSKI W., DEPA Ł., KANTURSKI M., WĘGIEREK M., WIECZOREK K. 2015. An annotated checklist of the aphids (Hemiptera: Aphidomorpha) of Poland. *Polish Journal of Entomology* 84(4): 383–420.

Accepted: 19 June 2019; published: 29 November 2019

Licensed under a Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>